

FIRST DETECTION OF NEAR-INFRARED INTRADAY VARIATIONS IN THE SEYFERT 1 NUCLEUS NGC4395

TAKEO MINEZAKI^{1,2}, YUZURU YOSHII^{1,3}, YUKIYASU KOBAYASHI⁴, KEIGO ENYA⁵, MASAHIRO SUGANUMA⁴, HIROYUKI TOMITA¹,
 SHINTARO KOSHIDA⁶, MASAHIRO YAMAUCHI⁶, AND TSUTOMU AOKI⁷

accepted for publication in *ApJL*, 04/07/2006

ABSTRACT

We carried out a one-night optical *V* and near-infrared *JHK* monitoring observation of the least luminous Seyfert 1 galaxy, NGC4395, on 2004 May 1, and detected for the first time the intraday flux variations in the *J* and *H* bands, while such variation was not clearly seen for the *K* band. The detected *J* and *H* variations are synchronized with the flux variation in the *V* band, which indicates that the intraday-variable component of near-infrared continuum emission of the NGC4395 nucleus is an extension of power-law continuum emission to the near-infrared and originates in an outer region of the central accretion disk. On the other hand, from our regular program of long-term optical *BVI* and near-infrared *JHK* monitoring observation of NGC4395 from 2004 February 12 until 2005 January 22, we found large flux variations in all the bands on time scales of days to months. The optical *BVI* variations are almost synchronized with each other, but not completely with the near-infrared *JHK* variations. The color temperature of the near-infrared variable component is estimated to be $T = 1320 \sim 1710$ K, in agreement with thermal emission from hot dust tori in active galactic nuclei (AGNs). We therefore conclude that the near-infrared variation consists of two components having different time scales, so that a small *K*-flux variation on a time scale of a few hours would possibly be veiled by large variation of thermal dust emission on a time scale of days.

Subject headings: galaxies: Seyfert — galaxies: active — infrared: galaxies — dust, extinction — galaxies: individual(NGC 4395)

1. INTRODUCTION

The near-infrared continuum emission of type 1 AGNs is considered to be dominated by thermal emission from hot dust surrounding the central engine (Kobayashi et al. 1993; Glikman, Helfand, & White 2006). However, the contribution of the accretion disk to the near-infrared continuum emission is suggested by the standard accretion disk model with large outer radius that predicts the power-law continuum emission of $f_\nu \propto \nu^{+1/3}$ at long wavelengths (Shakura & Sunyaev 1973). If this is the case, part of the near-infrared emission would originate in the outer accretion disk, although it is difficult to separate the accretion disk component from the dominant thermal dust component by the simple method of spectral decomposition. In order to clearly separate the accretion disk component, techniques using variability (Tomita et al. 2006) and polarization (Kishimoto, Antonucci, & Blaes 2005) were recently proposed and applied to some AGNs.

NGC4395 is a unique object known as the least luminous Seyfert 1 galaxy (Filippenko & Sargent 1989), having a bolometric luminosity of only $L_{\text{bol}} \sim 10^{40-41}$ ergs s^{-1} with broad emission lines in the UV and optical spectra (Filippenko & Sargent 1989; Filippenko, Ho, & Sargent

1993; Kraemer et al. 1999). Its spectral energy distribution (SED) from the X-ray to radio wavelengths resembles those of normal type 1 AGNs rather than those of low-ionization nuclear emission-line regions (LINERs) (Moran et al. 1999). Although a compact radio source is detected, it is a radio-quiet AGN (Ho & Ulvestad 2001; Wrobel, Fasnacht, & Ho 2001; Ho & Peng 2001). The mass of the central black hole measured by reverberation-mapping observation of the C IV emission line is as small as $M_{\text{BH}} = (3.6 \pm 1.1) \times 10^5 M_\odot$ (Peterson et al. 2005).

Moreover, the NGC4395 nucleus shows extreme variability at many wavelengths, probably because of its low luminosity and small black hole mass. Fast and large-amplitude variations were observed in the X-ray (Lira et al. 1999; Iwasawa et al. 2000; Shih, Iwasawa, & Fabian 2003; Vaughan et al. 2005; Moran et al. 2005), and intraday variations were observed in the UV and optical (Peterson et al. 2005; Skelton et al. 2005). In the near-infrared, while Quillen et al. (2000) reported a 15% flux variation in the *H* band at an interval of three weeks, no detailed intraday monitoring observation has been carried out yet.

In this paper, we report the results of the optical and near-infrared intraday monitoring observation of the Seyfert 1 nucleus in NGC4395. As observed in the X-ray, UV and optical, we found the intraday variation in the *J* and *H* bands almost synchronized with the optical intraday variation. We conclude that the emission of the intraday-variable component in the optical and near-infrared originates in the central accretion disk. Throughout this paper, we adopt 4.0 Mpc for the distance of NGC4395 (Thim et al. 2004).

2. OBSERVATIONS

A one-night monitoring observation was carried out on 2004/05/01 (UTC), while our regular program of long-term monitoring observation started from 2004/02/12 (UTC), using the multicolor imaging photometer (MIP) mounted on the

¹ Institute of Astronomy, School of Science, University of Tokyo, 2-21-1 Osawa, Mitaka, Tokyo 181-0015, Japan

² e-mail: minezaki@mtk.ia.s.u-tokyo.ac.jp

³ Research Center for the Early Universe, School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

⁴ National Astronomical Observatory, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

⁵ Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3-1-1, Yoshinodai, Sagami-hara, Kanagawa, 229-8510, Japan

⁶ Department of Astronomy, School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0013, Japan

⁷ Kiso Observatory, Institute of Astronomy, School of Science, University of Tokyo, 10762-30 Mitake, Kiso, Nagano 397-0101, Japan




FIG. 1.— The optical and near-infrared light curves of the NGC4395 nucleus on 2004/05/01 (UTC); V (small open circles), J (open squares), H (small filled circles), and K (open circles). For clarity, an offset of -0.05 mJy is applied to H, and -0.25 mJy to K.

2 m telescope of the Multicolor Active Galactic Nuclei Monitoring (MAGNUM) project at the Haleakala Observatories in Hawaii (Kobayashi et al. 1998a,b; Yoshii 2002). Here, we describe the procedures of observation and data reduction only briefly, because most of them have been described in Minezaki et al. (2004).

Our one-night monitoring observation started at 06:31 and ended at 12:35 on 2004/05/01 (UTC). Using the MIP's simultaneous optical and near-infrared imaging capability, a sequence of (V,K), (V,H), and (V,J) imagings was repeated cyclically during the observation, and 59 data points for V and 20 data points for each of J, H, and K were obtained. The average monitoring interval was six minutes for V and 18 minutes for J, H, and K. In addition, the long-term monitoring observation started from 2004/02/12 (UTC) and 11 data points were obtained until 2005/01/22 (UTC). After 2004/04/29 (UTC), the observation each night was carried out quasi-simultaneously (within 10 minutes) in the B, V, I, J, H, and K bands.

The nuclear flux was measured relative to a nearby reference star ($\alpha = 12^{\text{h}}25^{\text{m}}50.91^{\text{s}}$, $\delta = +33^{\circ}33'10.1''$ (J2000)), which was observed along with the target, within the field of view of the MIP. The optical fluxes of the star were calibrated with respect to a photometric standard star in Landolt (1992), and the near-infrared fluxes were taken from the 2MASS All-Sky Catalog of Point Sources (Cutri et al. 2003). The diameter of photometric aperture was 2.8 arcsec for the optical, and 2.4 arcsec for the near-infrared. The flux of the host galaxy within the aperture was not subtracted.

3. RESULTS

Figure 1 shows the optical and near-infrared light curves of the NGC4395 nucleus on 2004/05/01 (UTC). Galactic extinction was corrected for according to Schlegel, Finkbeiner, & Davis (1998). Clear flux variations in the V, J, and H bands are found during the monitoring period of six hours, and this is the first detection of intraday variation in the near-infrared for normal Seyfert 1 galaxies. The amplitude of variations was $0.05 - 0.09$ mag, however, the real amplitude for the NGC4395 nucleus must be larger, because non-variable fluxes such as the host galaxy flux and the narrow-line flux were not subtracted from the data. On the other hand, intraday variation is not clearly seen in the K band.

Apparently, the intraday variations in the V, J, and H bands




FIG. 2.— The CCF (dashed line) and the CCCD (solid line) between the V-band and the J+H combined light curves on 2004/05/01 (UTC). The CCF is peaked at 0.003 days and the CCF centroid is estimated as $\tau_{\text{cent}} = 0.005_{-0.006}^{+0.011}$ days or $7.2_{-8.6}^{+15.8}$ minutes from the CCCD.

are almost synchronized. In order to estimate a possible lag between the optical and near-infrared intraday variations, we applied a cross-correlation analysis. First, in order to increase the number of data points, the J-band and the H-band light curves were combined according to the linear regressions of the J- and H-band fluxes to the V-band flux obtained simultaneously during the one-night monitoring observation. Then a cross-correlation function (CCF) was computed based on the linear interpolation method (Gaskell & Peterson 1987; White & Peterson 1994). Since the number of V-band data points is larger and their photometric accuracy is better, the V-band light curve was interpolated. The lag of the CCF centroid τ_{cent} was adopted to represent the lag between two light curves. The error of τ_{cent} was estimated by a model-independent Monte Carlo simulation called the “flux randomization/random subset selection” (FR/RSS) method (Peterson et al. 2004). The cross-correlation centroid distribution (CCCD) was made based on the simulation of 10000 realizations. Figure 2 shows the CCF and the CCCD between the V-band and the combined near-infrared light curves on 2004/05/01 (UTC). The near-infrared lag time behind V is estimated from the CCCD as $\tau_{\text{cent}} = 0.005_{-0.006}^{+0.011}$ days, or $7.2_{-8.6}^{+15.8}$ minutes. The intraday variations in the near-infrared (J, H) are almost synchronized with that in the optical within an accuracy of ~ 10 minutes. The short time scale of variation (a few hours) and the synchronization to the optical variation (~ 10 minutes) suggest that the intraday-variable component of flux in the near-infrared originates in the central accretion disk. That is, it is an extension of optical power-law continuum emission to near-infrared wavelength.

Figure 3 shows the optical and near-infrared light curves of the long-term monitoring observation of the NGC4395 nucleus. The optical and near-infrared fluxes are clearly variable on time scales of days to months, and the apparent amplitude of variation during the monitoring period reaches about a few tenths to one magnitude. The optical flux variations in the B, V, and I bands are synchronized with each other, while the near-infrared flux variations, especially in the K band, are not completely synchronized with those in the optical.

Figure 4 shows the flux-flux diagrams from the long-term monitoring observation. The variations in the B or I bands relative to the V band are almost on straight lines, which suggests that the spectral shape of variable components in the optical region stays constant as reported for Seyfert galaxies (Winkler et al. 1992; Winkler 1997). The slope of these




FIG. 3.— The optical and near-infrared light curves of the NGC4395 nucleus; *B* (open squares), *V* (small filled circles), and *I* (open circles) in the lower panel, and *J* (open squares), *H* (small filled circles), and *K* (open circles) in the upper panel. An offset of 0.1 mJy is applied to *I* for clarity. The intraday variation data of 2004/05/01 (UTC) are presented by vertical bars at MJD= 53126. The optical data at the last two epochs were noisy because of instrumental problems, and their *B*-band data were omitted.




FIG. 4.— The flux-flux diagrams from the long-term monitoring of the NGC4395 nucleus. Plotted against *V* are the data for *B* (filled squares and crosses) and *I* (filled circles) in the left panel, and *J* (open squares), *H* (small filled circles), and *K* (open circles) in the right panel. The straight lines in the left panel are the regression lines based on the *BVI* data (filled symbols) which were taken almost simultaneously within 10 minutes.

lines, or the flux variation gradient (FVG), corresponds to the power-law index of variable component. From the regression analysis using the *B*, *V*, and *I* data points of the quasi-simultaneous observation, we obtain $\alpha_\nu = -0.6 \pm 0.2$ for *B* to *V*, and $\alpha_\nu = 0.0 \pm 0.1$ for *V* to *I*, which are similar to those of more luminous Seyfert galaxies (Winkler 1997).

In contrast, the near-infrared *J*, *H*, and *K* variations largely deviate from straight lines, and the deviation is larger for longer wavelengths, which suggests that another variable component exists in the near-infrared. We extracted the SED of this component by evaluating the near-infrared flux difference at two epochs for which the optical fluxes were almost the same. The flux difference between 2004/04/29 and

2004/07/21 is Δf (mJy) = 0.025 ± 0.009 (*B*), 0.009 ± 0.010 (*V*), 0.020 ± 0.006 (*I*), 0.071 ± 0.009 (*J*), 0.162 ± 0.011 (*H*), and 0.332 ± 0.015 (*K*). The color temperature assuming the black body radiation is estimated as 1710 ± 150 K from $\Delta f_H / \Delta f_J$, and 1320 ± 80 K from $\Delta f_K / \Delta f_H$. These estimates are consistent with the near-infrared SEDs of QSOs originating in the hot dust tori (Kobayashi et al. 1993; Glikman, Helfand, & White 2006).

4. DISCUSSION

The near-infrared intraday variation of the NGC4395 nucleus and its synchronization to the optical suggest that there exists a near-infrared emitting region in the accretion disk, so that the lag between the optical and the near-infrared intraday variations would be a direct measure of the extent of the accretion disk, if the reprocessing of incident flux variation to the near-infrared can be applied. Accordingly, a wavelength-dependent lag in variations of power-law continuum emission is expected, interpreted in terms of light-travel time from the central energy source to each radius of the accretion disk (Collier et al. 1999).

Therefore, the outer radius of the accretion disk indicated by the possible lag between the *V*-band and the near-infrared intraday variations is worthwhile comparing with the size of the broad emission-line region (BLR). Our measurement of near-infrared lag time behind *V* is $\tau_{\text{cent}} = 0.005^{+0.006}_{-0.011}$ days (2004/05/01). Peterson et al. (2005) reported from the reverberation mapping program for the C IV emission line of the NGC4395 nucleus that the C IV lag time behind UV continuum ($\lambda = 1350$ Å) is $\tau_{\text{CIV}} = 0.033^{+0.017}_{-0.013}$ days (2004/04/10) and $\tau_{\text{CIV}} = 0.046^{+0.017}_{-0.013}$ days (2004/07/03). Apparently, τ_{cent} is smaller than τ_{CIV} . If we adopt the wavelength-dependent relation $\tau(\lambda) \propto \lambda^{4/3}$ given by Collier et al. (1999), the near-infrared lag behind the UV continuum would be $\sim 1.3 - 1.4$ times larger than that behind *V*. In addition, the inner region of the BLR of the NGC4395 nucleus would be a few times smaller than the C IV emitting region because the BLR of Seyfert galaxies is stratified (Peterson & Wandel 1999). These considerations suggest that the outer radius of the accretion disk might be comparable to the inner radius of the BLR in the NGC4395 nucleus.

Next, we discuss the contribution of the power-law continuum component in the *K* band. As described in the previous section, thermal emission from the dust torus largely contributes to the *K*-band flux of the NGC4395 nucleus. Since the dust torus is expected to be extended, the short time scale variation of incident flux is smeared out and the thermal flux would not be variable on such a time scale. Then, even if there was an intraday flux variation of the power-law continuum component in the *K* band, it would be veiled by the large amount of thermal flux from the dust torus. We estimated an average of *K*-band fluxes at peaks (five data points) and at valleys (six data points) of the *V*-band intraday variation, and derived their difference. The 3σ upper limit of the flux difference of 0.04 mJy represents an amount of intraday variation in the *K* band. On the other hand, such a difference in the *J* and *H* bands is about 0.03 mJy. Therefore, it is possible to consider that an intraday flux variation in the *K* band is comparable to those in the *J* and *H* bands.

Finally, we estimate the inner radius of the dust torus according to the correlation between the *V*-band absolute magnitude and the lag time of *K*-band flux variation reported by Minezaki et al. (2004). The host galaxy flux within the

aperture is estimated as 0.26 mJy from the *I*-band magnitude of the central star cluster in NGC4395 (Filippenko & Ho 2003) and by assuming typical *V* – *I* color of Scd galaxies (Fukugita, Shimasaku, & Ichikawa 1995). The flux of the [O III] narrow line was taken from Skelton et al. (2005), and its contamination of the *V*-band flux is estimated as 0.24 mJy. After subtracting these fluxes, the *V*-band absolute magnitude of the NGC4395 nucleus is estimated as $M_V \sim -11$ mag, which leads to a lag time of $\tau_K \sim 1$ day. The thermal flux from such a dust torus, whose inner radius is ~ 1 light-day, would not be variable on a time scale of hours, veiling the possible intraday variation of the power-law continuum component in the *K* band.

The present paper indicates that more intense multi-wavelength monitoring observation of the NGC4395 nucleus every 10 minutes a night will definitely extract the synchronized intraday variation of power-law continuum component up to the near-infrared *K* band, and a wavelength-dependent lag behind X-ray or UV is able to probe the radial structure

of central accretion disk. This intense multi-wavelength monitoring observation over a month will alternatively extract the near-infrared *K* variation of thermal dust component, and its lag behind UV or optical gives an accurate measure of the inner radius of minimum dust torus in the least luminous nucleus among Seyfert 1 AGNs. These lag measurements of greater accuracy, together with the C IV lag (Peterson et al. 2005), will finally uncover a full range of radial structure consisting of the accretion disk, the BLR, and the dust torus in the NGC4395 nucleus.

We thank staff at the Haleakala Observatories for their help with facility maintenance. This research has been supported partly by the Grant-in-Aid of Scientific Research (10041110, 10304014, 11740120, 12640233, 14047206, 14253001, 14540223, 16740106) and COE Research (07CE2002) of the Ministry of Education, Science, Culture and Sports of Japan.

REFERENCES

- Collier, S., Horne, K., Wanders, I., & Peterson, B. M. 1999, MNRAS, 302, 24
- Cutri, R. M. et al. 2003, The 2MASS All-Sky Catalog of Point Sources
- Gaskell, C. M., & Peterson, B. M. 1987, ApJS, 65, 1
- Filippenko, A. V., & Sargent, W. L. W. 1989, ApJ, 342, L11
- Filippenko, A. V., Ho, L. C., & Sargent, W. L. W. 1993, ApJ, 410, L75
- Filippenko, A. V., & Ho, L. C. 2003, ApJ, 588, L13
- Fukugita, M., Shimasaku, K., & Ichikawa, T. 1995, PASP, 107, 945
- Glikman, E., Helfand, D. J., & White, R. L. 2006, ApJ accepted (astro-ph/0511640)
- Ho, L. C., & Peng, C. Y. 2001, ApJ, 555, 650
- Ho, L. C., & Ulvestad, J. S. 2001, ApJS, 133, 77
- Iwasawa, K., Fabian, A. C., Almaini, O., Lira, P., Lawrence, A., Hayashida, K., & Inoue, H. 2000, MNRAS, 318, 879
- Kobayashi, Y., Sato, S., Yamashita, T., Shiba, H., & Takami, H. 1993, ApJ, 404, 94
- Kobayashi, Y., Yoshii, Y., Peterson, B. A., Miyazaki, S., Aoki, T., Minezaki, T., Kawara, K., Enya, K., Okada, N., Suganuma, M., Green, B., O'Brien, M., & Lawrence, K. 1998a, Proc. SPIE Vol. 3352, pp120-128, Advanced Technology Optical/IR Telescopes VI, ed. L. M. Stepp
- Kobayashi, Y., Yoshii, Y., Peterson, B. A., Minezaki, T., Enya, K., Suganuma, M., & Yamamuro, T. 1998b, Proc. SPIE Vol. 3354, pp769-776, Infrared Astronomical Instrumentation, ed. A. M. Fowler
- Kishimoto, M., Antonucci, R., & Blaes, O. 2005, MNRAS, 364, 640
- Kraemer, S. B., Ho, L. C., Crenshaw, D. M., Shields, J. C., & Filippenko, A. V. 1999, ApJ, 520, 564
- Landolt, A. U. 1992, AJ, 104, 340
- Lira, P., Lawrence, A., O'Brien, P., Johnson, R. A., Terlevich, R., & Bannister, N. 1999, MNRAS, 305, 109
- Minezaki, T., Yoshii, Y., Kobayashi, Y., Enya, K., Suganuma, M., Tomita, H., Aoki, T., & Peterson, B. A. 2004, ApJ, 600, L35
- Moran, E. C., Filippenko, A. V., Ho, L. C., Shields, J. C., Belloni, T., Comastri, A., Snowden, S. L., & Sramek, R. A. 1999, PASP, 111, 801
- Moran, E. C., Eracleous, M., Leighly, K. M., Chartas, G., Filippenko, A. V., Ho, Luis C., & Blanco, P. R. 2005, AJ, 129, 2108
- Peterson, B. M., & Wandel, A. 1999, ApJ, 521, L95
- Peterson, B. M., Ferrarese, L., Gilbert, K. M., Kaspi, S. K., Malkan, M. A., Maoz, D., Merritt, D., Netzer, H., Onken, C. A., Pogge, R. W., Vestergaard, M., & Wandel, A. 2004, ApJ, 613, 682
- Peterson, B. M., Bentz, M. C., Desroches, L.-B., Filippenko, A. V., Ho, L. C., Kaspi, S., Laor, A., Maoz, D., Moran, E. C., Pogge, R. W., & Quillen, A. C. 2005, ApJ, 632, 799
- Quillen, A. C., Shaked, S., Alonso-Herrero, A., McDonald, C., Lee, A., Rieke, M. J., & Rieke, G. H. 2000, ApJ, 532, L17
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525
- Shakura, N. I., & Sunyaev, R. A. 1973, A&A, 24, 337
- Shih, D. C., Iwasawa, K., & Fabian, A. C. 2003, MNRAS, 341, 973
- Skelton, J. E., Lawrence, A., Pappa, A., Lira, P., & Almaini, O. 2005, MNRAS, 358, 781
- Tomita, H. et al. in preparation
- Thim, F., Hoessel, J. G., Saha, A., Claver, J., Dolphin, A., & Tammann, G. A. 2004, AJ, 127, 2322
- Vanden Berk, D. E. et al. 2001, AJ, 122, 549
- Vaughan, S., Iwasawa, K., Fabian, A. C., & Hayashida, K. 2005, MNRAS, 356, 524
- White, R. J. & Peterson, B. M. 1994, PASP, 106, 879
- Winkler, H., Glass, I. S., van Wyk, F., Marang, F., Jones, J. H. S., Buckley, D. A. H., & Sekiguchi, K. 1992, MNRAS, 257, 659
- Winkler, H. 1997, MNRAS, 292, 273
- Wrobel, J. M., Fassnacht, C. D., & Ho, L. C. 2001, ApJ, 553, L23
- Yoshii, Y. 2002, in New Trends in Theoretical and Observational Cosmology, ed. K. Sato and T. Shiromizu (Tokyo: Universal Academy Press), 235